



Contribution ID: 513

Type: Oral 20 Minutes

## Viscous fingering and nonlinear waves in a Langmuir adsorbed solute

*Tuesday, 15 May 2018 16:16 (15 minutes)*

Waves in chromatography are well known to the practitioners. Characteristics of these waves are strongly correlated to the nature of the adsorption isotherm. We model the displacement of a finite slice containing an adsorbed solute by a carrier liquid flow. We examine the nonlinear dynamics that emerge from the interactions of rarefaction, shock layer and/or viscous fingers (VF) in the finite solute slice that adsorbs on a porous matrix according to a Langmuir isotherm. The differences between a linear and a Langmuir isotherm are discussed. In the absence of VF, a shock layer (rarefaction) wave appears at the frontal (rear) interface of the solute. VF at a viscously unstable rarefaction interface propagates through the finite sample to preempt the shock layer. However, no such incident is observed when the shock layer front features VF. Various quantities (e.g., the onset of VF, shock layer thickness, etc.) are calculated as a function of the saturation rate and the results are supported by mathematical analysis.

### References

### Acceptance of Terms and Conditions

[Click here to agree](#)

**Primary author:** RANA, Chinar

**Co-authors:** Prof. MARTIN, Michel (ESPCI-Paris, France); Dr PRAMANIK, Satyajit (NORDITA, Stockholm, Sweden); Dr MISHRA, Manoranjan (Indian Institute of Technology Ropar); DE WIT, Anne (Universite libre de Bruxelles (ULB))

**Presenter:** RANA, Chinar

**Session Classification:** Parallel 5-A

**Track Classification:** MS 2.05: Modeling and Controlling of Viscous Fingering in Miscible and Immiscible Displacements in Subsurface Porous Media