



Contribution ID: 379

Type: Oral 20 Minutes

Modification of wettability and interfacial tension by biosurfactant-producing bacteria for geologic carbon storage

Monday, 14 May 2018 11:18 (15 minutes)

Injection of carbon dioxide (CO₂) into deep geologic formations has been widely proposed as an effective way for the permanent storage of CO₂. Modification of the interfacial properties of CO₂ in minerals by using surfactant has been proposed aiming on increasing the mobility of CO₂ through porous media. Surfactants are proven to effectively alter the interfacial tension and wettability in both CO₂/water/mineral system, improving the displacement and sweep efficiencies of CO₂ in porous media. In the meantime, biosurfactants have been drawing much attention as an alternative to the chemical surfactants for their biodegradability, ecological suitability and low toxicity. However, the question as to the extent of microbial alterations in fluid wettability and interfacial tension under reservoir pressure and temperature conditions still warrants further investigation. Therefore, this study investigated the role of lipopeptide biosurfactant on wettability and interfacial tension alterations in a CO₂/brine/mineral system for different CO₂ phases during the growth of thermotolerant and barotolerant bacteria, *Bacillus subtilis*, and the production of lipopeptide biosurfactant, surfactin. Quartz, mica and carbonate substrates were selected and used as representative minerals. While monitoring the changes in the interfacial tension and wettability with pH, fluid samples were acquired from the brine phase, and the concentrations of glucose, nitrate, ammonium and surfactin in the acquired samples were quantitatively assessed using various assays and spectroscopic methods. As a result of surfactin production by *B. subtilis*, we observed the reductions in interfacial tension and increases in contact angle at all tested cases. The concentration of surfactin and the rate of wettability alteration differed with the experimental conditions. The modification of CO₂ wettability was the greatest for liquid CO₂ while the least of modification was observed for gaseous CO₂. The obtained results allow in-depth assessment of the feasibility of using biosurfactant-producing bacteria for effective geologic carbon storage practices.

References

Park, T., Joo, H.W., Kim, K.Y., Kim, S., Yoon, S., and Kwon, T.H. (2017) "Biosurfactant as an enhancer of geologic carbon storage: Microbial modification of interfacial tension and contact angle in carbon dioxide/water/quartz systems", *Frontiers in Microbiology*

Acceptance of Terms and Conditions

[Click here to agree](#)

Primary authors: Mr PARK, Taehyung (Korea Advanced Institute of Science and Technology (KAIST)); Ms JEON, Minkyung (Korea Advanced Institute of Science and Technology (KAIST)); Prof. KIM, Seunghee (University of Nebraska–Lincoln); Prof. KWON, Tae-Hyuk (Korea Advanced Institute of Science and Technology (KAIST))

Presenter: Mr PARK, Taehyung (Korea Advanced Institute of Science and Technology (KAIST))

Session Classification: Parallel 1-A

Track Classification: MS 3.11: fundamental aspects of geological storage of CO₂