InterPore 10th Annual Meeting and Jubilee
May 14-17 2018, New Orleans, USA
interpore.org/neworleans

Contribution ID: **905**   Type: **Poster + 3 Minute Pitch**

Wicking as partially-saturated flow of a liquid in thin swelling porous media

Monday 14 May 2018 15:52 (2 minutes)

Unsaturated flow through thin layers of porous media are encountered in many industrial applications, including the liquid-absorbing hygiene products such as wipes, paper towels, and diapers [1-4]. These consumer products demand specific absorbent properties with storage of liquid playing a significant role. Understanding fluid flow and deformation processes in thin swelling porous media is critical for the development and design of these products.

We use an averaging approach [5-17] for modeling a system consisting of multiple layers of thin, absorbing swelling porous media as the layer-wise 2D interacting continua, to rigorously derive a 2D averaged macroscopic mass-balance model for each layer and to develop the required constitutive relationships for a system of thin porous layers made of one liquid (water) and two deformable solid phases (fiber and hydrogel). The developed model consists of a set of partial differential equations that keep track the time dependent behavior of variables such as piezometric head, saturation, porosity, and layer thickness, as the liquid moves throughout the multi-layered porous medium. Hence, this model can be used to describe the absorbency process [18], to predict and understand the flow and storage of a liquid in conjunction with the deformation of layers in multilayered thin porous media that is absorbing the liquid, and swelling during deformation. This model will enormously improve the computational speeds, allowing one to develop a fast and reasonably accurate simulation of the unsaturated flow.

The numerical simulations are carried out with the flow parameters and geometries for a few representative cases such as wicking into dry horizontal and inclined porous plates. The simulation predictions, which predict detailed 2-D flow fields, are found to be in good agreement with the experimental and 3D computational results.

References

1. Buchholz, F.L., Model of liquid permeability in swollen composites of superabsorbent polymer and fiber. Journal of applied polymer science, 2006. 102(4): p. 4075-4084.

2. Buchholz, F.L. and A.T. Graham, Modern superabsorbent polymer technology. John! Wiley & Sons, Inc, 605 Third Ave, New York, NY 10016, USA, 1998. 279, 1998.

3. Diersch, H.-J.G., et al., Modeling unsaturated flow in absorbent swelling porous media: Part 2. Numerical simulation. Transport in Porous Media, 2011. 86(3): p. 753-776.

4. Diersch, H.-J.G., et al., Modeling unsaturated flow in absorbent swelling porous media: Part 1. Theory. Transport in Porous Media, 2010. 83(3): p. 437-464.

5. Majid Hassanizadeh, W.G.G., General conservation equations for multi-phase systems: 1. Averaging procedure. Advances in Water Resources, 1979. 2: p. 131-144.

6. Hassanizadeh, S.M., Derivation of basic equations of mass transport in porous media, Part 2. Generalized Darcy's and Fick's laws. Advances in Water Resources, 1986. 9(4): p. 207-222.

7. Hassanizadeh, M. and W. Gray, General conservation equations for multi-phase system: 2. Mass, momenta, energy and entropy equations. Adv. Water Resour, 1979. 2(3): p. 191-203.

8. Preziosi, L. and A. Farina, On Darcy's law for growing porous media. International Journal of Non-Linear Mechanics, 2002. 37(3): p. 485-491.

9. Majid Hassanizadeh, W.G.G., General conservation equations for multi-phase systems: 3. Constitutive theory for porous media flow. 1980. 3(1): p. 25-40.

10. Gray, W., et al., Mathematical tools for changing spatial scales in the analysis of physical systemsCRC Press. Boca Raton, 1993.

11. Hassanizadeh, S.M. and W.G. Gray, Mechanics and thermodynamics of multiphase flow in porous media including interphase boundaries. Advances in water resources, 1990. 13(4): p. 169-186.

12. Gray, W.G., Constitutive theory for vertically averaged equations describing steam-water flow in porous media. Water Resources Research, 1983. 19(6): p. 1501-1510.

13. Gray, W.G., Derivation of vertically averaged equations describing multiphase flow in porous media. Water Resources Research, 1982. 18(6): p. 1705-1712.

14. Gray, W.G. and P. Lee, On the theorems for local volume averaging of multiphase systems. International Journal of Multiphase Flow, 1977. 3(4): p. 333-340.

15. Whitaker, S., The method of volume averaging. Vol. 13. 1998: Springer Science & Business Media.

16. Qin, C. and S. Hassanizadeh, A new approach to modelling water flooding in a polymer electrolyte fuel cell. international journal of hydrogen energy, 2015. 40(8): p. 3348-3358.

17. Qin, C. and S. Hassanizadeh, Multiphase flow through multilayers of thin porous media: General balance equations and constitutive relationships for a solid–gas–liquid three-phase system. International Journal of Heat and Mass Transfer, 2014. 70: p. 693-708.

18. Feldkamp, J.R., A mathematical description of liquid flow through partially saturated deformable porous media. report 2.02(TL No 7439.).

Acceptance of Terms and Conditions

Click here to agree

Authors: Dr KAFFEL, Ahmed; Prof. PILLAI, Krishna (University of Wisconsin Milwaukee)

Presenters: Dr KAFFEL, Ahmed; Prof. PILLAI, Krishna (University of Wisconsin Milwaukee)

Session Classification: Parallel 2-F

Track Classification: MS 4.14: Wicking of Liquids in Porous Materials