



Contribution ID: 444

Type: **Poster (+) Presentation**

Experimental Investigation of Capillary Number's Control on Stress-Dependent Shifts in Irreducible Saturation in Deformable Porous Media

Monday, 31 May 2021 19:35 (1 hour)

Characterization of irreducible saturation of the wetting phase during multiphase fluid flow in porous media is essential for an accurate estimation of CO₂ storage capacity and hydrocarbon recovery of the geological formations. Despite pore deformation has been shown to significantly control single-phase and multiphase fluid flow in porous media, the interactive controls of capillary number and mechanical pore deformation on irreducible saturation during multiphase fluid flow in geo-materials is not yet fully explored. In this study, the stress-dependent shifts of irreducible water saturation (S_{wir}) of a Berea sandstone and an Indiana limestone specimen are investigated through series of two-phase (water-N₂) core-flooding experiments (i.e., drainage) under increasing effective stress from 10 MPa to 30 MPa and isothermal (40°C) conditions. We used X-ray computed micro-tomography to quantify changes in the topology of the pore-space with effective stress. The controls of the capillary number on the stress-dependent shifts of S_{wir} is studied through experiments under constant injection rate and constant injection pressure conditions, independently. We find a 22% and 52% decrease in S_{wir} of Berea sandstone and Indiana limestone, respectively, in response to an increase in effective stress under constant injection rate (i.e., increasing capillary number) condition. We further find a 27% increase in S_{wir} of Indiana limestone with the same increase in effective stress under constant injection pressure (i.e., decreasing capillary number) condition. We reveal that the deformation of the pore throats, due to an excess effective confining stress, and changes in the driving energy for the gas phase to invade smaller channels, due to an increase/decrease in capillary number, leads to a decrease/increase in S_{wir} of both specimens. These micro-scale and macro-scale observations underscore the remarkable control of capillary number on deformation-dependent fluid-fluid displacement in porous media, which pave the way for relevant research in geoscience and engineering.

Time Block Preference

Time Block C (18:00-21:00 CET)

References

- Haghi, A.H., Chalaturnyk, R., Blunt, M.J., Hodder, K., Geiger, S., (2020). Poromechanical Controls on Multiphase Fluid Flow in Earth Material. *Scientific Reports*, 11, 3328.
- Haghi, A.H., Talman, S., Chalaturnyk, R. (2020). Consecutive Experimental Determination of Stress-dependent fluid flow Properties of Berea Sandstone and Implications for Two-phase Flow Modelling. *Water Resources Research*, 56(1)
- Haghi, A.H., Chalaturnyk, R., Talman, S. (2019) Stress-Dependent Pore Deformation Effects on Multiphase Flow Properties of Porous Media. *Scientific Reports*, 9(1), 1-10

Acceptance of Terms and Conditions

[Click here to agree](#)

Newsletter

Student Poster Award

Primary author: HAGHI, Amir (University of Alberta)

Co-author: CHALATURNYK, Richard (University of Alberta)

Presenter: HAGHI, Amir (University of Alberta)

Session Classification: Poster +

Track Classification: (MS4) Swelling and shrinking porous media