



Contribution ID: 643

Type: Oral Presentation

# **Evolution of Bentheimer Sandstone Wettability During Cyclic scCO**<sub>2</sub>-Brine Injections

Thursday, 3 June 2021 11:15 (15 minutes)

Geologic sequestration in sedimentary formations has been identified as a potential technology to prevent climate-change inducing carbon dioxide (CO2) from being emitted to the atmosphere. In order to achieve safe and effective storage underground, accurate understanding and predictions of CO<sub>2</sub> behavior in subsurface rock formations is required; including quantifying how much CO2 is trapped within pore spaces by capillarity (vs. how much remains mobile), and constraining the occurrence of physio-chemical reactions between CO<sub>2</sub> and the mineral matrix. Experiments where multiple cycles of CO<sub>2</sub> and brine are injected into rock samples have produced conflicting results; likely due to differences in mineral content of samples, pressure-temperature conditions, the suite of aqueous chemistry parameters which may impact results, and experimental apparatus and protocol. We present a new set of experiments, designed to replicate the conditions of a previous study, but with a new experimental design, apparatus, and timeline. We confirm the previous results that demonstrated shifts in CO<sub>2</sub> trapping behaviour over multiple injection cycles, and we conduct additional analyses to discern the fluid-fluid macroscopic contact angle, curvature, interfacial area, and topology of trapped CO2 ganglia. The results are considered in the context of two mechanisms that could cause evolution of  $CO_2$  behavior; (1) mineral fines migration, and (2) solubility-induced  $CO_2$  adhesion and wettability alteration. We do not observe significant evidence of fines migration in this "clean" Bentheimer sandstone system. Conversely, Lattice-Boltzmann simulations based on the proposed conceptual model of wettability alteration exhibit similar trends in pressure measurments, end-point saturation levels, and CO<sub>2</sub> ganglia morphological characteristics as experiments, indicating that the changes observed in experiments could be due to wettability alteration as proposed. We observe that this wettability alteration renders  $CO_2$ more stable in the rock pore space, increasing capillary trapping over four injection cycles.

### **Time Block Preference**

Time Block A (09:00-12:00 CET)

#### References

## Acceptance of Terms and Conditions

Click here to agree

Newsletter

#### **Student Poster Award**

Primary author: HERRING, Anna (The Australian National University)

**Co-authors:** LI, Zhe (Advanced Research Computing, Virginia Polytechnic Institute & State University, Blacksburg, Virginia); SUN, Chenhao (University of New South Wales); ARMSTRONG, Ryan; MCCLURE, James (Virginia Tech); SAADATFAR, Mohammad (Australian National University (ANU))

Presenter: HERRING, Anna (The Australian National University)

Session Classification: MS1

Track Classification: (MS1) Porous Media for a Green World: Energy & Climate