



Contribution ID: 189

Type: Oral Presentation

# **A Unified Plot for Fluid Phase Transitions**

Thursday, 3 June 2021 19:00 (15 minutes)

While many mesoporous materials are found naturally (clays, coal, and shale), many others are synthetized for industrial applications like sensing, fluid separations, and energy storage. For the optimization of these materials for various applications, an accurate determination and understanding of the pore size distribution and morphology is vital.

Fluid phase transitions employ the alteration of physical properties of confined fluids as markers for pore space characterization. This is expressly revealed by the Kelvin equation for gas-liquid transitions and the Gibbs-Thomson equation for solid-liquid transitions. As may be thought, for one and the same material, both methods are expected to yield the same pore size distribution with little or no deviations. However, it has been shown that, both approaches may yield slightly different results.

Herein, we introduce a unified plot for both gas-liquid and solid-liquid transitions based on the Kelvin and Gibbs-Thomson equations. We analyze three mesoporous silica solids; SBA-15, Vycor and MCM-41, in order of decreasing pore size. Remarkable similarity is observed in the fluid phase transitions in the SBA-15 material of pore size approximately 9nm. With decreasing pore size, a divergence between both transitions is observed. We postulate that, this may primarily be due to the increasing influence of thermodynamic fluctuations with decreasing pore size.

## **Time Block Preference**

Time Block B (14:00-17:00 CET)

#### References

[1] Schneider D.; Kondrashova D.; Valiullin R., 2017, "Phase transitions in disordered mesoporous solids", Scientific Reports, 7, 7216.

[2] Schneider, D. and Valiullin R., 2019, "Capillary condensation and evaporation in irregular channels: Sorption isotherm for serially connected pore model", Journal of Physical Chemistry C, 123, 16239.

[3] Enninful H.R.N.B., Schneider D., Hoppe A., König S., Fröba M., Enke D. and Valiullin R., 2019, "Comparative gas sorption and cryoporometry study of mesoporous glass structure: Application of the serially connected pore model", Frontiers in Chemistry, doi: 10.3389/fchem.2019.00230.

[4] Enninful H.R.N.B., Schneider D., Kohns R., Enke D. and Valiullin R., 2020, "A novel approach for advanced thermoporometry characterization of mesoporous solids: Transition kernels and the serially connected pore model", Microporous and Mesoporous Materials 309, 110534.

[5] Enninful H.R.N.B., Schneider D., Enke D. and Valiullin R., 2021, "Impact of Geometrical Disorder on Phase Equilibria of Fluids and Solids Confined in Mesoporous Materials", Langmuir, Under Revision.

#### Acceptance of Terms and Conditions

Click here to agree

### Newsletter

## **Student Poster Award**

Yes, I would like to enter this submission into the student poster award

**Primary authors:** ENNINFUL, Henry (Universität Leipzig); ENKE, Dirk (Leipzig University); VALIULLIN, Rustem (Leizpig University)

Presenter: ENNINFUL, Henry (Universität Leipzig)

Session Classification: MS13

Track Classification: (MS13) Fluids in Nanoporous Media