

**HYDRUS and analytical modeling of seepage in porous banks of commingled
ephemeral streams having triangular flash-flood hydrographs: emergence and
extinction of an “ephemeral” unconfined aquifer**

A. R. Kacimov¹, and J. Šimůnek²

¹Department of Soils, Water and Agricultural Engineering, Sultan Qaboos University, Oman

ORCID ID: orcid.org/0000-0003-2543-3219

Emails: anvar@squ.edu.om, akacimov@gmail.com

²Department of Environmental Sciences, University of California Riverside, CA, USA

Email: jsimunek@ucr.edu

Short title for running head: Emergence and extinction of an “ephemeral” unconfined aquifer

Submitted to minisymposium MS25 “Subsurface Water Flow and Contaminant Transport
Processes – Special Session in Honour of Harry Vereecken”

Interpore

<https://events.interpore.org/event/25/>.

on ..., 2021 as abstract N...,
first set of referees comments received on ...;
revised version submitted: on;
accepted:.... rejected:

Address for correspondence:

Prof. Kacimov A.R., Department of Soils, Water and Agricultural Engineering,
College of Agricultural and Marine Sciences, Sultan Qaboos University,
Al-Khod 123, PO Box 34 Sultanate of Oman

Fax (968) 24413-418

Emails: anvar@squ.edu.om

akacimov@gmail.com

Omani page:

<https://www.squ.edu.om/agriculture/Academic-Department/Soils-Water-and-Agricultural-Engineering>

Abstract

Perched aquifers (PA) are important sources of groundwater in arid/semiarid climates where surface water resources are irregular (see, e.g., *Intermittent...*, 2017). PAs as hydrostratigraphic units are commonly conceived as being created/maintained by vertical infiltration from a vadose zone above aquifer's phreatic surface and vanished due to a) vertical percolation, which descends into a subjacent vadose zone under aquifer's thin low-permeable bed, and b) vertical ascending seepage flow driven by evapotranspiration. In our work, we study "ephemeral" aquifers, whose advent and disappearance are controlled by prevalently horizontal fluxes, one into a vadose zone and another into a conterminous ephemeral stream. Initial-boundary value problems to the Richards 2-D equation, 1-D Boussinesq equation, and 2-D Laplace equation are solved in models of saturated-unsaturated and purely saturated flows. At $t < 0$, both the stream and adjacent bank (wedge-shaped in a vertical cross-section) are dry. At $t > 0$, a flash flood takes place. Wadi's water first imbibes into the bank and later exfiltrates back through the slope, which is made of two discharging segments: a seepage face and ponded interface between stream's water and the aquifer (Fig. 1a).




Fig. 1. a) A vertical cross-section of wadi at three phases of a flash flood and a conjugated "ephemeral" aquifer; b) Contour map of the pore pressure head during the drawup phase.

52

53 The process is controlled by a specified wadi hydrograph. We focus on the case of a constant rate
 54 drawup-drawdown regime, i.e., the wadi stage rises-drops linearly with time. Analytical
 55 (Barenblatt's and Barenblatt's type, see Barenblatt et al., 1984) and numerical (HYDRUS-2D, see
 56 Šimůnek et al., 2016) solutions compare well. Dynamics of bank's phreatic surface (conjugated
 57 with the wadi water level), transient fluxes, pore-water storage in the bank, loci of the tips of the
 58 wetting fronts (propagating along a bedrock of an emerged "ephemeral" unconfined aquifer), pore
 59 pressure isobars (in particular, the position of the crest of an evolving groundwater mound),
 60 piezometric contours, vector fields of Darcian velocity, isotachs, and streamlines in the three
 61 models are studied. Fig. 1b illustrates a snapshot of HYDRUS pore pressure heads for the drawup
 62 phase. A rapid drawup of the wadi level and slow post-flash-flood drawdown (i.e., a triangular
 63 "synthetic" hydrograph) generate an intricate topology of groundwater and soil moisture motion,
 64 viz. the three phases of expansion-slumping-evanescence (E-S-E). For example, in the slumping
 65 phase, a stagnation point emerges on the bedrock at a certain time and flow bifurcates into one
 66 halve, which keeps moving to the right in Fig. 1a, and another discharges through the slope.

67 The mound may or may not be intercepted by a dormant (not-pumping) well in the riparian zone.
 68 The well is tracked by HYDRUS observational nodes. Depending on the phase of aquifer's
 69 evolution ("yes"- "yes"- "no" in the example of Fig. 1b), the distance between the well and wadi,
 70 screen's depth, and hydraulic properties of the porous medium, the phreatic surface either intersects
 71 the well or not. In the former case, a well operator can start abstracting groundwater. Implications
 72 for intelligent interception of short-lived groundwater pockets by on-bank farms and for
 73 groundwater banking in MAR projects are discussed. Another application of our HYDRUS and
 74 analytical models is in the ecohydrology of riparian phreatophytes (e.g., the Christ-thorn trees in
 75 Oman, see, e.g., Al-Maktoumi et al., 2020). Ecotones of this wild vegetation are aligned with

wadis' and can serve as bioindicators of the post-flash-flood extension of ephemeral aquifers in Fig. 1a.

References

- +Al-Maktoumi, A., Kacimov, A., Al-Busaidi, H., Al-Ismaïly, S., Al-Mayahi, A., Al-Khanbashi, S., and Al-Sulaimi, A. (2020). Enhancement of infiltration rate of clogged porous beds in the vicinity of dams in arid zones by the roots of indigenous *Ziziphus spina-christ* trees. *Hydrological Processes*, 34(22), 4226-4238.
- +Barenblatt, G. I., Entov, V. M., and Ryzhik, V. M., 1990. *Theory of Fluid Flows Through Natural Rocks*. Kluwer, Dordrecht.
- +Datry, T., N. Bonada, and A. Boulton (eds.), 2017. *Intermittent Rivers and Ephemeral Streams*. Elsevier, Cambridge, Massachusetts.
- +Šimůnek, J., van Genuchten, M. Th., and Šejna, M., 2016. Recent developments and applications of the HYDRUS computer software packages, *Vadose Zone J.*, 15(7), pp. 25, doi: 10.2136/vzj2016.04.0033.

User Name: anvarkacimov
Password: EFcQnD8Z