



Contribution ID: 423

Type: Oral Presentation

Engineering biofilm hydraulic resistance on the microscale

Tuesday, 31 May 2022 17:30 (15 minutes)

Bacterial biofilms are amongst the most successful modes of life in the terrestrial environment and ubiquitous within porous systems, such as soils and membrane interfaces (1). The bacteria within a biofilm are bound together by self-secreted extracellular polymeric substances (EPS), yielding a natural gel-like structure (2). EPS provides a protective shield and structural architecture to constituent cells of the biofilm, which shapes internal mass transport within the biofilm and flow field of the colonized porous media (3). The interplay between biofilm formation and flow leads to the formation of preferential flow pathways and significant increases in membrane hydraulic resistance, as investigated in macroscale experimental systems (4, 5). However, the complex multiphase interaction between biofilm proliferation and flow within irregular porous media make the deduction of scalable physical relationships challenging. Consequently, a mechanistic understanding linking EPS composition, biofilm morphogenesis and porosity with hydraulic resistance is still missing (6).

In this work, we investigate the time evolution of biofilm morphology and hydraulic resistance as a function of flow conditions and EPS composition. We grow the model bacterium *Bacillus subtilis* in two types of microfluidic channel and use a library of EPS mutants devoid of polysaccharide (EPS-O), protein (TasA) or hydrophobin (BslA) to modulate EPS composition. Firstly, we grow and characterize the biofilm during its morphogenesis in a novel microfluidic platform where the biofilm is grown on a cellulose nanofibril (CNF) membrane barrier. The CNF platform allowed precise measurement of single colony biofilm hydraulic resistance as a function of pressure driven flow and EPS composition, whilst retaining full optical access. To validate the scalability of our findings, we use an unconfined microfluidic geometry, a model 2D porous media, which increases the biofilm volume by approximately two orders of magnitude. We measure hydraulic resistance and apply advanced optical visualization techniques to quantify biofilm morphogenesis and internal transport. This study allowed us to quantify how flow and EPS composition shape the hydraulic resistance of biofilm at the micro and meso scale. The relationships derived from these studies enhance our understanding of how we can engineer biofilms to tune their hydraulic resistance.

Acceptance of the Terms & Conditions

[Click here to agree](#)

MDPI Energies Student Poster Award

No, do not submit my presentation for the student posters award.

Country

Switzerland

References

1. H.-C. Flemming, et al., Biofilms: an emergent form of bacterial life. *Nat Rev Micro* 14, 563–575 (2016).

2. P. Di Martino, Extracellular polymeric substances, a key element in understanding biofilm phenotype. *AIMS Microbiol.* 4, 274–288 (2018).
3. M. Peszynska, A. Trykozko, G. Iltis, S. Schlueter, D. Wildenschild, Biofilm growth in porous media: Experiments, computational modeling at the porescale, and upscaling. *Adv. Water Resour.* 95, 288–301 (2016).
4. N. Hassannayebi, et al., Relationship Between Microbial Growth and Hydraulic Properties at the Sub-Pore Scale. *Transp. Porous Media* 139, 579–593 (2021).
5. M. Carrel, et al., Pore-Scale Hydrodynamics in a Progressively Bioclogged Three-Dimensional Porous Medium: 3-D Particle Tracking Experiments and Stochastic Transport Modeling. *Water Resour. Res.* 54, 2183–2198 (2018).
6. P. Desmond, et al., Controlling the hydraulic resistance of membrane biofilms by engineering biofilm physical structure. *Water Res.* 210, 118031 (2022).

Time Block Preference

Time Block A (09:00-12:00 CET)

Participation

In person

Primary authors: SECCHI, Eleonora (ETH Zürich); KURZ, Dorothee Luise; Mr BOGGON, Cameron (ETH Zurich); CHARLTON, Sam (ETH Zurich)

Presenter: CHARLTON, Sam (ETH Zurich)

Session Classification: MS05

Track Classification: (MS05) Biochemical processes and biofilms in porous media