



Contribution ID: 250

Type: Oral Presentation

## Flow-direction dependence of upscaled capillary pressure-saturation curve

Monday, 30 May 2022 14:25 (15 minutes)

Immiscible two-phase flow is widely present in natural and synthetic processes. The flow behaviour of two fluids is governed by constitutive relations, relative permeability and capillary pressure. These empirical relations are often influenced by the dynamic of the process and the characteristics of porous media such as heterogeneity. The effect of the non-equilibrium condition on the capillary pressure-saturation behaviour has been investigated and shown that the dynamic capillary pressure is different from the one measure under the equilibrium condition. Moreover, recent studies showed that the presence of micro-heterogeneity in porous media changed the trend and the extent of the capillary pressure-saturation compared to the background porous medium. Although, the magnitude and the trend of the capillary pressure-saturation curves remained almost unchanged irrespective of the direction of the fluid flow in these studies.

The present work investigates the effect of heterogeneity interface on saturation distribution and capillary pressure-saturation behaviour in a micromodel study. The micro model is made up of two sections called fine and coarse sections. Microfluidic experiments and optical imaging and analyses were used to calculate capillary pressure and saturation of fluids. Drainage experiments were conducted at four different flow rates with a wide range of capillary numbers in both directions (i.e. fine to coarse and coarse to fine). The saturation of each phase was measured using image analysis. Moreover, the capillary pressure at the pore scale was calculated by estimating the curvature of each fluid-fluid interface. Then using the fluid-fluid interfacial surface area, the averaged capillary pressure in the coarse section, fine section and the entire micromodel, was calculated.

Results show that the averaged dynamic capillary pressure-saturation curve with the presence of a heterogeneity interface does not follow the monotonic shape of the conventional capillary pressure curve, measured under equilibrium conditions. Moreover, the results demonstrate a non-monotonic relationship between the remaining wetting phase saturation and the capillary number. It is mainly due to the competition between the capillary and viscous forces during the transition from capillary fingering to viscous fingering regime. The results reveal that considering the flow direction with respect to the heterogeneity interface, can lead to a better prediction of the upscaled capillary pressure-saturation relation and the remaining wetting phase saturation.

### Acceptance of the Terms & Conditions

[Click here to agree](#)

### MDPI Energies Student Poster Award

Yes, I would like to submit this presentation into the student poster award.

### Country

United Kingdom

## References

- An, S. a.-B. (2020). Transition From Viscous Fingering to Capillary Fingering: Application of GPU-Based Fully Implicit Dynamic Pore Network Modeling. *Water Resources Research*, e2020WR028149.
- Das, D. B.-A. (2004). A numerical study of micro-heterogeneity effects on upscaled properties of two-phase flow in porous media. *Transport in Porous Media*, 329–350.
- Dawe, R. A. (2011). Microscale visual study of end effects at permeability discontinuities. *Transport in porous media*, 601–616.
- Joekar-Niasar, V. a. (2012). Uniqueness of specific interfacial area–capillary pressure–saturation relationship under non-equilibrium conditions in two-phase porous media flow. *Transport in Porous Media*, 465–486.
- Lenormand, R. (1986). Pattern growth and fluid displacements through porous media. *Physica A: Statistical Mechanics and its Applications*, 114–123.
- Rabbani, H. S.-Y. (2018). Suppressing viscous fingering in structured porous media. *PNAS*, 4833-4838.
- van Duijn, C. J. (1995). The effect of capillary forces on immiscible two-phase flow in heterogeneous porous media. *Transport in Porous Media*, 71–93.

## Time Block Preference

Time Block B (14:00-17:00 CET)

## Participation

Online

**Primary author:** SHOKRI, Javad (University of Manchester)

**Co-authors:** GODINEZ BRIZUELA, Omar Emmanuel (The University of Manchester); ERFANI , Hamidreza (Norwegian University of Science and Technology); Mr CHEN, Yongqiang (University of Manchester); BABAEI, Masoud (University of Manchester); BERKOWITZ, Brian (Weizmann Institute of Science); NIASAR, Vahid (University of Manchester)

**Presenter:** NIASAR, Vahid (University of Manchester)

**Session Classification:** MS11

**Track Classification:** (MS11) Microfluidics and nanofluidics in porous systems