InterPore2023



Contribution ID: 986 Type: Oral Presentation

Magnetic Resonance Measurements of Fluids in Shale

Tuesday, 23 May 2023 15:40 (30 minutes)

Shale formations are of increasing importance world-wide as petroleum reservoirs. Conventional core analysis measurements are ill suited to the analysis of shale samples since the pore fluids present are very difficult to extract due to the very small pore sizes. Magnetic Resonance (MR) permits non-invasive analysis of intact samples and is therefore an intuitively appealing method for shale analysis. The MR relaxation correlation measurement T1-T2 is now commonly employed for analysis of shale fluids. The T1-T2 measurement struggles to observe and quantify the fast decaying signal from pore fluids in shales. A new method, which we term T1-T2captures the very short-lived fluid shale MR signal exceptionally well and provides well resolved discrimination and quantification of pore fluids, as well as the solid-like kerogen. In this lecture we will review MR relaxation time measurements of fluids in porous media, consider the broad category of MR relaxation correlation measurements ultimately leading to T1-T2 analysis of shales.

Participation

In-Person

References

MDPI Energies Student Poster Award

No, do not submit my presenation for the student posters award.

Country

Canada

Acceptance of the Terms & Conditions

Click here to agree

Energy Transition Focused Abstracts

Primary author: BALCOM, Bruce (University of New Brunswick)

Presenter: BALCOM, Bruce (University of New Brunswick)

Track Classification: (MS24) Invited and Plenary Sessions