



Contribution ID: 939

Type: Oral Presentation

Thermodynamic properties of ganglia in heterogeneous porous media

Monday, 22 May 2023 11:35 (15 minutes)

Ganglia (bubbles, or droplets) are widespread in porous media of various industrial applications. Thermodynamic properties of a ganglion, such as its morphology, free energy, capillary pressure, surface energy, etc., are crucial in determining its transport and reactive performance. Although these in homogeneous porous media have been recently resolved [1, 2], it is still challenging to quantitatively describe the thermodynamic properties of ganglia in heterogeneous media [3-5].

We develop a pore-scale numerical algorithm for determining the thermodynamic properties of hydrostatic ganglia in heterogeneous porous media. We track cycles of quasi-static growth and quasi-static shrinkage of a ganglion in a two-dimensional heterogeneous porous media, as shown in Figure.1(a). The algorithm is as follows:

- (1) Create a heterogeneous porous medium and set the initial capillary pressure (P_c) and pore occupancy of the ganglion.
- (2) Find the hydrostatic morphology of the ganglia with set P_c and pore occupancy, and record its properties including ganglion volume (V), free energy (F), surface area (A), etc.
- (3) Make a small change in P_c while keeping pore occupancy unchanged, and then check whether a new hydrostatic morphology can be achieved. If so, the change is reversible and we go back to step (2) and continue. If not, an irreversible event emerges that changes pore occupancy while keeping V unchanged, and we search for the new stable morphology.

This algorithm may be used to simulate degassing and dissolution process in heterogeneous porous media and enhance our understanding of these processes. In addition, this algorithm may be used to construct the energy landscape of the entire heterogeneous porous media. We believe that this work helps better understand the behaviors of the dispersed phase in heterogeneous porous media.




Figure 1: (a) Snapshots of ganglion growth in heterogeneous porous media. (b) Evolution of the capillary pressure, surface free energy, and specific surface area with ganglion volume during ganglion growth in heterogeneous porous media.

Participation

In-Person

References

1. Armstrong, R.T., et al., Porous Media Characterization Using Minkowski Functionals: Theories, Applications and Future Directions. *Transport in Porous Media*, 2018. 130(1): p. 305-335.
2. Wang, C., Y. Mehmani, and K. Xu, Capillary equilibrium of bubbles in porous media. *Proceedings of the National Academy of Sciences*, 2021. 118(17): p. e2024069118.
3. Li, Y., C. Garing, and S.M. Benson, A continuum-scale representation of Ostwald ripening in heterogeneous porous media. *Journal of Fluid Mechanics*, 2020. 889.
4. Lin, Q., et al., Drainage Capillary Pressure Distribution and Fluid Displacement in a Heterogeneous Laminated Sandstone. *Geophysical Research Letters*, 2021. 48(14).
5. Krishnamurthy, P.G., D. DiCarlo, and T. Meckel, Geologic Heterogeneity Controls on Trapping and Migration of CO₂. *Geophysical Research Letters*, 2022. 49(16).

MDPI Energies Student Poster Award

No, do not submit my presentation for the student posters award.

Country

China

Acceptance of the Terms & Conditions

[Click here to agree](#)

Energy Transition Focused Abstracts

Primary authors: WANG, Chuanxi (Peking University); XU, Ke (Peking University)

Presenter: WANG, Chuanxi (Peking University)

Session Classification: MS09

Track Classification: (MS09) Pore-scale modelling