InterPore2023



Contribution ID: 1021

Type: Poster Presentation

Asymptotic Homogenization - Modeling the charging behavior of Li-ion batteries

Wednesday, 24 May 2023 16:10 (1h 30m)

The modeling of lithium-ion cells plays an important role in the electrification of many industries, such as the automotive industry. Models with high accuracy and low computational complexity are essential for the optimization of such applications.

To achieve this, a fully homogenized macroscale (FHM) model has been proposed [1]. This reduced model is based on effective mass and charge transport equations in isothermal active particles and electrolyte.

In current-limited battery operation, and thus at low C-rates, the FHM model works very well. At high Crates, the predictions of the FHM model deviate significantly from those of the Direct Numerical Simulations (DNS). A high C-rate leads to high current densities between the active particles and the electrolyte and thus to steep concentration gradients near the interface. The resulting deviation between the inner concentration and surface concentration can no longer be covered by the homogenized value. Consequently, the FHM model is not applicable to diffusion-limited battery operation.

Therefore, we propose an extension of the FHM model based on Wang's diffusion length concept for spheres [2]. Here, the diffusion length is an additional variable that characterizes the correlation between the homogenized concentration and the surface concentration as a function of the C-rate.

This extension allows to consider the diffusion limitation within the active particles, thus allowing a reliable application at higher C-rates.

The extended FHM model is validated using a direct numerical simulation (DNS) for the charge of a graphite half-cell. Despite the non-spherical properties of the graphite electrode, the diffusion length concept can be successfully applied. The concentrations and potential distributions of the FHM model are in good agreement with the DNS results.

Participation

In-Person

References

[1] Arunachalam, H., & Onori, S. (2019). Full homogenized macroscale model and pseudo-2-dimensional model for lithium-ion battery dynamics: comparative analysis, experimental verification and sensitivity analysis. Journal of The Electrochemical Society, 166(8), A1380 https://doi.org/10.1149/2.0051908jes

[2] Wang, C. Y., Gu, W. B., & Liaw, B. Y. (1998). Micro-Macroscopic Coupled Modeling of Batteries and Fuel Cells: I. Model Development. Journal of the Electrochemical Society, 145(10), 3407. https://doi.org/10.1149/1.1838820

MDPI Energies Student Poster Award

Yes, I would like to submit this presentation into the student poster award.

Country

Germany

Acceptance of the Terms & Conditions

Click here to agree

Energy Transition Focused Abstracts

Primary author: Mr LOMBARDO, Giuliano (University of Stuttgart, Institute of Chemical Process Engineering)

Co-authors: Mr YUANZHEN, Wang (Max Planck Institute for Medical Research, Department of Cellular Biophysics); Prof. NIEKEN, Ulrich (Institute of Chemical Process Engineering, University of Stuttgart)

Presenter: Mr LOMBARDO, Giuliano (University of Stuttgart, Institute of Chemical Process Engineering)

Session Classification: Poster

Track Classification: (MS19) Elastic, electrical, and electrochemical processes and properties in porous media