



Contribution ID: 261

Type: Oral Presentation

Pore aperture regulated surface adsorption and mass transfer of hydrocarbon and CO₂ in organic nanopores

Monday, 13 May 2024 13:25 (15 minutes)

Shale oil, widely distributed in organic (i.e., kerogen) nanopores, is playing an ever-increasing role in addressing the global energy crisis, but is faced with challenges of low recovery efficiency due to well-developed nanopores. It is believed that the pore size distribution of kerogen is in the range of several angstroms to tens of nanometers (AAPG bulletin 96 (6): 1099-1119). In such a narrow pore space, oil molecules are dominated by adsorbed phase, which is hard to recover relying on pressure drop (International Journal of Coal Geology 147 (2015): 9-24). CO₂ huff-and-huff is identified as a promising method to enhance oil recovery while achieving CO₂ sequestration. Clarifying the adsorption and extraction behaviors of hydrocarbons in kerogen nanopores is crucial for accurately predicting oil recovery and revealing CO₂ enhanced oil recovery (CO₂-EOR) mechanisms.

In this work, we adopted molecular dynamics (MD) simulation to study the static spatial distribution and dynamic mass transfer of hydrocarbon and CO₂ in slit-shaped kerogen nanopores by carefully designing a series of pore apertures. It shows that the adsorption and extraction behaviors of oil molecules are closely related to pore aperture. Interestingly, we found that the surface adsorption of oil molecules demonstrates a non-monotonic trend of rising after falling as the pore size decreases. Specifically, when the pore size is reduced to a certain value, oil molecules exhibit a pseudo-double layer adsorption state, in which the surface adsorption peaks of oil molecules are significantly weakened. On the other hand, although the reduction of pore width adversely affects the extraction speed of oil during CO₂ soaking, the recovery efficiency presents a jump for the oil at pseudo-double layer adsorption state. Meanwhile, the surface adsorption of CO₂ is also greatly enhanced, which leads to the highest CO₂/Oil ratio in the nanopores. Collectively, our work provides fresh and important insights into hydrocarbon occurrence state and CO₂-EOR mechanisms in organic-rich shale reservoirs and builds up a good foundation for accurate predictions of oil recovery.

Acceptance of the Terms & Conditions

[Click here to agree](#)

Student Awards

I would like to submit this presentation into both awards

Country

China

Porous Media & Biology Focused Abstracts

This abstract is related to Porous Media & Biology

References

Characterization of gas shale pore systems by porosimetry, pycnometry, surface area, and field emission scanning electron microscopy/transmission electron microscopy image analyses: Examples from the Barnett, Woodford, Haynesville, Marcellus, and Doig units. AAPG bulletin 96 (6): 1099-1119; Oil adsorption in shale nanopores and its effect on recoverable oil-in-place. International Journal of Coal Geology 147 (2015): 9-24.

Conference Proceedings

I am not interested in having my paper published in the proceedings

Primary author: ZHANG, mingshan (东北大学)

Presenter: ZHANG, mingshan (东北大学)

Session Classification: MS13

Track Classification: (MS13) Fluids in Nanoporous Media