



Contribution ID: 668

Type: Oral Presentation

A combined ionic Lewis acid descriptor and machine-learning approach to prediction of efficient oxygen reduction electrodes for ceramic fuel cells

Tuesday, 14 May 2024 15:00 (15 minutes)

Improved, highly active cathode materials are needed to promote the commercialization of ceramic fuel cell technology. However, the conventional trial-and-error process of material design, characterization and testing can make for a long and complex research cycle. Here we demonstrate an experimentally validated machine-learning-driven approach to accelerate the discovery of efficient oxygen reduction electrodes, where the ionic Lewis acid strength (ISA) is introduced as an effective physical descriptor for the oxygen reduction reaction activity of perovskite oxides. Four oxides, screened from 6,871 distinct perovskite compositions, are successfully synthesized and confirmed to have superior activity metrics. Experimental characterization reveals that decreased A-site and increased B-site ISAs in perovskite oxides considerably improve the surface exchange kinetics. Theoretical calculations indicate such improved activity is mainly attributed to the shift of electron pairs caused by polarization distribution of ISAs at sites A and B, which greatly reduces oxygen vacancy formation energy and migration barrier.

F

Acceptance of the Terms & Conditions

[Click here to agree](#)

Student Awards

Country

China

Porous Media & Biology Focused Abstracts

This abstract is related to Porous Media & Biology

References

Conference Proceedings

I am not interested in having my paper published in the proceedings

Primary author: ZHAI, Shuo (Shenzhen University)

Presenter: ZHAI, Shuo (Shenzhen University)

Session Classification: MS22

Track Classification: (MS22) Manufactured Porous Materials for Industrial Applications