



Contribution ID: 320

Type: **Poster Presentation**

Investigation on pore structure and imbibition characteristic of tight sandstone by nuclear magnetic resonance

Wednesday, 15 May 2024 09:25 (1h 30m)

Abstract: Tight sandstone reservoir, as unconventional oil and gas resource with abundant quantity, plays a significant role in supporting the growing global demand of energy consumption. However, tight sandstone has complex pore structure and strong heterogeneity, which brings challenges to the development of unconventional hydrocarbon resource. In recent years, the nuclear magnetic resonance method has been widely applied in porous media with the advantages of nondestructive, fast and wider range of pore size characterization in investigating pore structure. But pore size distribution curve cannot be obtained directly by nuclear magnetic resonance method. The transverse relaxation time spectrum from nuclear magnetic resonance could be converted into the pore size distribution curve by appropriate surface relaxivity, which can be affected by paramagnetic minerals and varies with lithology. Using low temperature nitrogen adsorption method, high-speed centrifugation method, and nuclear magnetic resonance, the surface relaxivity of five tight sandstone samples from different depths of Dongying depression was measured in this study. And then the influence of mineral composition from experimental samples on surface relaxivity was discussed. In addition, the pore size distribution curve from mercury intrusion porosimetry is utilized to verify the accuracy of the pore size distribution curve inverted by the surface relaxivity. Based on the obtained surface relaxivity, imbibition experiments under high temperature and high pressure conditions were conducted combined with nuclear magnetic resonance technique. Then the flow characteristics of whole cores and different pore size types (micropores, mesopores and macropores) influenced by temperature and pressure was analyzed, and the imbibition mechanism of tight sandstones resource was revealed.

Keywords: NMR, Surface relaxivity, Pore size distribution, Imbibition

Acceptance of the Terms & Conditions

[Click here to agree](#)

Student Awards

Country

China

Porous Media & Biology Focused Abstracts

References

Reference 1. Coates, G.R., Xiao, L., and Prammer, M.G., NMR Logging: Principles and Applications, Houston: Haliburton Energy Services, 1999. 2. Kleinberg, R.L., Kenyon, W.E., and Mitra, P.P., Mechanism of NMR Relaxation of Fluids in Rock, J. Magn. Reson. Series A, vol. 108, no. 2, pp. 206-214, 1994. 3. Lyu, C., Ning, Z., and Cole, D.R., Experimental Investigation on T2 Cutoffs of Tight Sandstones: Comparisons between Outcrop and Reservoir Cores, J. Petrol. Sci. Eng., vol. 191, 2020. 4. Sigal, R.F., Pore Size Distributions for Organic Shale Reservoir Rocks from Nuclear Magnetic Resonance Spectra Combined with Adsorption Measurements, SPE J., vol. 20, pp. 824-830, 2015

Conference Proceedings

I am interested in having my paper published in the proceedings.

Primary author: XIA, Xuanzhe (China University of Petroleum-Beijing)

Co-author: Prof. CAI, Jianchao (China University of Petroleum (Beijing))

Presenter: XIA, Xuanzhe (China University of Petroleum-Beijing)

Session Classification: Poster

Track Classification: (MS18) Innovative Methods for Characterization, Monitoring, and In-Situ Remediation of Contaminated Soils and Aquifers