



Contribution ID: 965

Type: Oral Presentation

Orthogonally different mineral reactions, same outcome of permeability reduction: How can this be?

Sustainable energy technologies that involve subsurface gas storage require reliable containment of buoyant fluids. An example is geologic carbon sequestration in which large volumes of CO₂ are injected deep underground into porous formations with overlying caprocks. Storage security could be jeopardized if fractures exist, so strategies are needed to seal permeable flow paths. In our work, two orthogonally different mineral reaction scenarios were explored. In one case minerals precipitated and in the other case minerals dissolved, but both cases had the same outcome of reduced fracture permeability. How can this be? In the first case, vein minerals from a mudrock sample of the Wolfcamp formation provided insights about syntaxial mineral growth in a fracture. Dolomite and other carbonate minerals had precipitated in the fracture, closing it off to fluid flow. In the second case, a carbonate-rich shale was reacted leading to calcite dissolution along fracture surfaces. Subsequent compression from normal stress collapsed the altered layer, sealing the fracture and reducing permeability. These studies show that multiple mineral reaction mechanisms can achieve fracture sealing and permeability reduction, a favorable outcome in subsurface applications where the goal is to reduce leakage risks.

Acceptance of the Terms & Conditions

[Click here to agree](#)

Student Awards

Country

USA

Porous Media & Biology Focused Abstracts

References

Conference Proceedings

I am not interested in having my paper published in the proceedings

Primary author: PETERS, Catherine

Presenter: PETERS, Catherine

Session Classification: Plenary/Invited

Track Classification: (MS25) Invited & Plenary Speakers